

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE New Records of Anopheles belenrae Rueda (Diptera: Culicidae) in North Korea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Walter Reed Biosystematics Unit, Silver Spring, MD, 20910				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

NOTE

New Records of *Anopheles belenrae* Rueda (Diptera: Culicidae) in North Korea

Anopheles belenrae Rueda is one of about 30 species of the Hyrcanus Group. This group contains about three-quarters of the species that comprise the Myzorrhynchus Series of genus *Anopheles* Meigen subgenus *Anopheles* in the Oriental and Palearctic Regions (Rueda et al. 2006). We report new records of *An. belenrae* from two provinces in North Korea (Democratic People's Republic of Korea). Adult specimens of this species were collected from Hwanghaebuk Province: Undok Dong (38°11'N, 126°40'E), Kaesong City, 10 August 2004, Coll. # NK027, 1 female; Songnam Dong (38°7'N, 126°38'E), Kaesong City, 9 August 2004, Coll. # NK028, 1 female; and Pyongyannam Province: Jansan Ri, Pyongsong City (38°53'N, 125°3'E), 15 July 2004, Coll. # NK025, 033, 2 females. Two species of the Hyrcanus Group, *An. sinensis* and *An. pullus*, also were found in the two provinces. Adults of *An. sinensis* were collected in Hwanghaebuk Province: Undok Dong, Kaesong City, 10 August 2004, Coll. # NK014-016, 3 females; Songnam Dong, Kaesong City, 9 August 2004, Coll. # NK017-021, NK029-031, 8 females; and Pyongyannam Province: Jansan Ri, Pyongsong City, 15 July 2004, Coll. # NK023-033, 036, 3 females. Adults of *An. pullus* were collected from Hwanghaebuk Province: Undok Dong, Kaesong City, 10 August 2004, Coll. # NK001, 002, 2 females; and Pyongyannam Province: Jansan Ri, Pyongsong City, 15–17 July 2004, Coll. # NK 003, 004-012, 12 females; 27–28 July 2004, Coll. # K022-023, 3 females. Adults of these three species were collected using aspirators from inside human dwellings or animal barns and outdoors, particularly in rice growing areas. Specimens

were identified morphologically (Rueda 2005, Tanaka et al. 1979) and verified by comparing our rDNA internal transcribed spacer 2 (ITS2) sequences to Wilkerson et al. (2003) and GenBank sequences (e.g., ITS2 Acc. No. AY375466, *An. belenrae*; AY375464, *An. sinensis*; AY375471, *An. pullus*).

Anopheles belenrae was first described from specimens collected in 2001 from Gyeonggi Province, South Korea, specifically Tongilchon (37°51'N, 126°47'E). It also occurs in Ogam-ri, Paju (37°49'N, 126°43'E), Gangwha, Incheon (37°45'N, 126°29'E), and Ogeum-ri, Thanhymyeon, Paju-si (37°49'N, 126°43'E) (Rueda 2005, Rueda et al. 2006). In 2006, Dr. Terry Klein and members of the 8th Medical Command, US Army, South Korea (in litt.) collected additional specimens of this species from the provinces of Gyeonggi, Chungcheongbuk and Gangwon. Adults were usually caught resting in cattle sheds of dairy farms, between 2000 and 2,230 h. The larvae were collected from various habitats including rice and paddy fields, ditches, marshes, marsh depressions, seepage-spring, ponds, stream margins, stream pools, rock pools, and storage tank pits (T. Klein, personal communication). In mainland China, *An. belenrae* occurs in the provinces of Liaoning and Shandong (Rueda et al. 2007). As in South Korea and North Korea, the status of this species as a malaria vector in China is unknown. Furthermore, the larval habitats of this species in North Korea and China are unknown, but they may be similar to those found in South Korea.

Many thanks to Y. Linton, B. Khuntirat, and C. Li for help with the

molecular analysis of the mosquito samples. Special thanks also to Y.-M. Huang, C. R. Summers, and B. P. Rueda for reviewing the manuscript. This research was performed under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense.

LITERATURE CITED

- Rueda, L. M. 2005. Two new species of *Anopheles* (*Anopheles*) *Hyrcanus* Group (Diptera: Culicidae) from the Republic of South Korea. *Zootaxa* 941: 1-26.
- Rueda, L. M., H. C. Kim, T. Klein, J. Pecor, C. Li, R. Sithiprasasna, M. Debboun, and R. C. Wilkerson. 2006. Distribution and larval habitat characteristics of *Anopheles* *Hyrcanus* Group and related mosquito species (Diptera: Culicidae) in South Korea. *Journal of Vector Ecology* 31: 199-206.
- Rueda, L. M., T. Zhao, Y. J. Ma, Q. Gao, Z. Guo Ding, B. Khuntirat, J. Sattabongkot, and R. C. Wilkerson. 2007. Updated distribution records of the *Anopheles* (*Anopheles*) *hyrcanus* species-group (Diptera: Culicidae) in China. *Zootaxa* 1407: 43-55.
- Tanaka, K., K. Mizusawa, and E. Saugstad. 1979. A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contributions of the American Entomological Institute* (Ann Arbor) 16: 1-987.
- Wilkerson, R. C., C. Li, L. M. Rueda, H. C. Kim, T. A. Klein, G. H. Song, and D. Strickman. 2003. Molecular confirmation of *Anopheles* (*Anopheles*) *lesteri* from the Republic of South Korea and its genetic identity with *An. (Ano.) anthropophagus* from China (Diptera: Culicidae). *Zootaxa* 378: 1-14.
- Leopoldo M. Rueda and Qi Gao, (LMR), *Walter Reed Biosystematics Unit, WRAIR, MSC MRC 534, Smithsonian Institution, 4210 Silver Hill Road, Suitland, MD 20746, U.S.A. (e-mail: ruedapol@si.edu); (QG) Jiangsu Institute of Parasitic Diseases, Meiyuan, Wuxi, Jiangsu 214064, P.R. China*